

Publisher's Note: "Polymer-specific effects of bulk relaxation and stringlike correlated motion in the dynamics of a supercooled polymer melt" [J. Chem. Phys. 119, 5290 (2003)]

M. Aichele

*Institut Charles Sadron, 6 rue Boussingault, 67083 Strasbourg, France
and Institut für Physik, Johannes Gutenberg-Universität, 55099 Mainz, Germany*

Y. Gebremichael

*Department of Chemical Engineering, University of Michigan, Ann Arbor, Michigan 48109
and Chemical Physics Program, Institute for Physical Sciences and Technology, University of Maryland,
College Park, Maryland 20742*

F. W. Starr^{a)}

*Center for Theoretical and Computational Materials Science and Polymers Division,
National Institute of Standards and Technology, Gaithersburg, Maryland 20899*

J. Baschnagle^{b)}

Institut Charles Sadron, 6 rue Boussingault, 67083 Strasbourg, France

S. C. Glotzer^{c)}

*Departments of Chemical Engineering and Materials Science and Engineering, University of Michigan,
Ann Arbor, Michigan 48109*

[DOI: 10.1063/1.1690246]

This article was originally published with an incorrect version of Fig. 8 on p. 5298. AIP apologizes for this error; the correct version of Fig. 8 appears below. All online versions of the article have been corrected.




FIG. 8. Temperature dependence of the ratio of $\langle s_{\text{seg}}(t_{\text{str}}^{\text{max}}) \rangle$ and $\langle s(t_{\text{str}}^{\text{max}}) \rangle$. $t_{\text{str}}^{\text{max}}$ is the peak time of $\langle s_{\text{seg}} \rangle$ and $\langle s \rangle$ at different temperatures. $T_{\text{MCT}} = 0.45$.

^{a)} Present address: Department of Physics, Wesleyan University, Middletown, CT 06459.

^{b)} Author to whom correspondence should be addressed. Electronic mail: baschnag@ics.u-strasbg.fr

^{c)} Author to whom correspondence should be addressed. Electronic mail: sglotzer@umich.edu